

Case Study 1

1. What is the MAX pharyngeal residue with liquids?

- a. <10%
- b. 10-49%
- c. 50-90%
- d. >90%

2. What is the MAX pharyngeal residue with pudding?

- a. <10%
- b. 10-49%
- c. 50-90%
- d. >90%

3. What is the MAX pharyngeal residue with cracker?

- a. <10%
- b. 10-49%
- c. 50-90%
- d. >90%
- e. consistency not given

4. What is the Pattern of Pharyngeal Residue?

- a. All bolus types presented
- b. Any bolus type (liquid, pudding, and/or cracker/cookie)
- c. Solids only (cracker and/or cookie)
- d. Liquids and/or pudding
- e. Any (but not all) bolus types presented

5. What is the Efficiency Grade?

- a. E0
- b. E1
- c. E2
- d. E3
- e. E4

Case Study 2

1. What is the MAX pharyngeal residue with liquids?

- a. <10%
- b. 10-49%
- c. 50-90%
- d. >90%

2. What is the MAX pharyngeal residue with pudding?

- a. <10%
- b. 10-49%
- c. 50-90%
- d. >90%

3. What is the MAX pharyngeal residue with cracker?

- a. <10%
- b. 10-49%
- c. 50-90%
- d. >90%
- e. consistency not given

4. What is the Pattern of Pharyngeal Residue?

- a. All bolus types presented
- b. Any bolus type (liquid, pudding, and/or cracker/cookie)
- c. Solids only (cracker and/or cookie)
- d. Liquids and/or pudding
- e. Any (but not all) bolus types presented

5. What is the Efficiency Grade?

- a. E0
- b. E1
- c. E2
- d. E3
- e. E4

Case Study 3

1. What is the MAX pharyngeal residue with liquids?

- a. <10%
- b. 10-49%
- c. 50-90%
- d. >90%

2. What is the MAX pharyngeal residue with pudding?

- a. <10%
- b. 10-49%
- c. 50-90%
- d. >90%

3. What is the MAX pharyngeal residue with cracker?

- a. <10%
- b. 10-49%
- c. 50-90%
- d. >90%
- e. consistency not given

4. What is the Pattern of Pharyngeal Residue?

- a. All bolus types presented
- b. Any bolus type (liquid, pudding, and/or cracker/cookie)
- c. Solids only (cracker and/or cookie)
- d. Liquids and/or pudding
- e. Any (but not all) bolus types presented

5. What is the Efficiency Grade?

- a. E0
- b. E1
- c. E2
- d. E3
- e. E4